

228/16

**GOAT EMBRYO PRODUCTION FROM *IN VITRO*
MATURED HETEROGENEOUS OOCYTES
FERTILISED BY INTRACYTOPLASMIC
SPERM INJECTION (ICSI) TECHNIQUE**

ABU NASAR MD. AMINOOR RAHMAN

**THESIS SUBMITTED IN FULFILMENT OF THE
REQUIREMENTS FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY**

**BIOHEALTH SCIENCE DIVISION
INSTITUTE OF BIOLOGICAL SCIENCES
FACULTY OF SCIENCE
UNIVERSITY OF MALAYA
KUALA LUMPUR, MALAYSIA**

FEBRUARY 2008


UNIVERSITY OF MALAYA

ORIGINAL LITERARY WORK DECLARATION

Name of Candidate : Abu Nasar Md. Aminoor Rahman
Passport No : Z 0826564 (New); N 0100569 (Old)
Registration No : SHB040003
Name of Degree : Doctor of Philosophy
Title of Thesis : Goat embryo production from *in vitro* matured heterogeneous oocytes fertilised by intracytoplasmic sperm injection (ICSI) technique
Field of Study : Reproductive Biotechnology

I do solemnly and sincerely declare that:


- (1) I am the sole author/writer of this Work;
- (2) This Work is original;
- (3) Any use of any work in which copyright exists was done by way of fair dealing and for permitted purposes and any excerpt or extract from, or reference to or reproduction of any copyright work has been disclosed expressly and sufficiently and the title of the Work and its authorship has been acknowledged in this Work;
- (4) I do not have any actual knowledge nor do I ought reasonably to know that the making of this Work constitutes an infringement of any copyright work;
- (5) I hereby assign all and every rights in the copyright to this Work to the University of Malaya ("UM"), who henceforth shall be owner of the copyright in this Work and that any reproduction or use in any form or by any means whatsoever is prohibited without the written consent of UM having been first had and obtained;
- (6) I am fully aware that if in the course of making this Work I have infringed any copyright whether internationally or otherwise, I may be subject to legal action or any other action as may be determined by UM.



Candidate's Signature

16.06.2008
Date

Subscribed and solemnly declared before,



Witness's Signature

16 June 2008
Date

Name : Prof Dr Ramli Abdullah
Designation : Institute of Biological Sciences
University of Malaya
50603 Kuala Lumpur

ABSTRACT

This study was conducted to produce goat embryos from heterogeneous oocytes retrieved by laparoscopic ovum pick-up (LOPU), ovariectomy and slicing of abattoir ovaries fertilised by intracytoplasmic sperm injection (ICSI) technique. For LOPU technique, selected does were oestrus synchronised and then superstimulated. Recovered oocytes were then cultured in microdrops of *in vitro* maturation (IVM) medium in presence of CO₂ (5%) at 38.5°C for 27 hours. Frozen buck semen was thawed and motile sperm were selected by swim-up. ICSI was performed on the lid of a culture dish (35 mm) under an inverted microscope fitted with micromanipulators. After catching and immobilising a motile sperm, it was then aspirated into the injection needle. Each mature oocyte was positioned in the ICSI microdrop with its first polar body at 6 or 12 o'clock on the holding pipette; and sperm was injected, head-first, into the ooplasm. Injected oocytes were then cultured for 7 to 8 days in mSOF medium in the CO₂ incubator. A sample of embryos from each stage and uncleaved oocytes were stained with Hoechst 33342 to examine the number of nuclei of the embryos or pronuclear status. The data were presented as mean±SEM and were analysed using one-way ANOVA. The significant differences among treatments were further analysed by DMRT and P<0.05 was considered significant.

In Experiment 1, two protocols, *i.e.*, P-1 and P-2 were compared. While 54.1% fertilisation and 29.8% cleavage rates were obtained with P-2, only 28.6% fertilisation rate without any cleavage was achieved with P-1. Some embryos with P-2 progressed to morula (4.0%); and this is a first report of goat embryo development by ICSI in Malaysia.

Experiment 2 evaluated the effect of time intervals (36, 60 and 72 hours) between FSH+hCG treatment and LOPU on oocyte recovery (OR) as well as IVM and ICSI outcomes. Significantly higher OR (P<0.001), IVM (P<0.001), fertilisation (P<0.05), cleavage (P<0.05) and morula development (P<0.05) rates were obtained with 60 and 72 hours interval groups than 36 hours. No pregnancy was achieved after embryo transfer (ET) at morula stage to recipient does.

In Experiment 3, the effect of calcium ionophore (Ca²⁺ ionophore) on oocyte activation and embryo development by ICSI was evaluated. A significantly (P<0.001) higher fertilisation (88.3%), cleavage (81.3%) and morula (32.5%) development rates were obtained with pre- & post-ICSI activation regimen than pre-ICSI, post-ICSI or ICSI control and all four sham (without sperm) injection groups. Staining of uncleaved oocytes and embryos revealed significantly (P<0.001) higher rates of 2 PN (47.5%) and oocyte activation (82.5%) with pre- & post-ICSI regimen than other groups.

Experiment 4 examined the effect of oocyte grading on IVM and embryo development by ICSI. Significantly higher ($P < 0.001$) maturation (95.2%) and morula development (73.3%), and higher ($P < 0.05$) fertilisation (74.2%) and cleavage (32.7%) rates were obtained from Grade A oocytes than Grades B, C and D. Similar results were obtained significantly when different grades of oocytes were treated with Ca^{2+} ionophore (pre- & post-ICSI regimen). However, no embryos progressed to blastocyst; and no pregnancy was achieved after ET at morula stage to recipient does.

In Experiment 5, developmental competence of oocytes of IVM and embryo of IVC from dysmorphic and normal groups were compared. Significantly ($P < 0.05$) higher maturation rates were obtained from oocytes having normal ooplasm (73.7%) than paler (54.6%) and big fat globules (56.3%); while no significant difference with darker ooplasm (69.0%). After ICSI, no significant difference was found between dysmorphic and normal oocytes in fertilisation and cleavage rates. However, dysmorphic oocytes were developmentally less competent than normal ones as most of these oocytes were either arrested or degenerated at around 5 to 8-cell stage; whereas, morula (12.6%) were obtained from normal oocytes. This is the first report of goat embryo development by ICSI from dysmorphic oocytes.

Experiment 6 evaluated the effect of oocyte source on OR, IVM and embryo developmental competence. LOPU technique provided better quality oocytes as evidenced by the recovery of higher proportion of Grade A oocytes (32.6%) than abattoir (19.0%) source. IVM rate was significantly ($P < 0.01$) higher in LOPU (73.8%) than abattoir (54.0%) source. Following ICSI, higher ($P < 0.05$) fertilisation rate, cleavage and morula were produced from LOPU than abattoir source.

It can be concluded from the present study that goat embryos could be produced through ICSI technique in Malaysia. Prolonging time interval between FSH+hCG treatment and LOPU from 36 to 60 and 72 hours could improve OR rate, oocyte quality, IVM and embryo development rates. Pre- & post-ICSI Ca^{2+} ionophore treatment regimen improves oocyte activation, fertilisation and embryo development rates. Grade A oocytes are more developmentally competent than other grades. Dysmorphic oocytes could also be matured and developed to embryos but are less competent than normal oocytes. LOPU can provide higher quality oocytes, better maturation and embryo development rates than abattoir source. These findings would be able to fill up some gaps of scientific information in goat ICSI. Therefore, considering the current goat population of Malaysia, ICSI coupled with LOPU technique would be an appropriate assisted reproductive technology (ART) to generate *in vitro* goat embryos which will, in turn, improve country's goat population.

ACKNOWLEDGEMENTS

First of all I would like thank to Almighty Allah for successfully completing my PhD research in good health.

I would like to express my cordial respect and thanks to my supervisor, Professor Dr. Ramli Bin Abdullah and co-supervisor Professor Dr. Wan Khadijah Wan Embong. Firstly, I want to show my gratefulness to Professor Dr. Ramli Bin Abdullah for giving me a chance to undertake the PhD study in the University of Malaya. I also thank him for his expert guidance and patience during the course of my experiments and thesis writing. I want to thank Professor Dr. Wan Khadijah Wan Embong for her scientific advice, enthusiasm, criticism and encouragements during the course of my study and thesis writing. I also acknowledge both of my supervisor's tremendous help during laparoscopic ovum pick-up and embryo transfer.

There are many people to thank for providing me help and guidance. In particular, Professor Ng Soon Chye, Adjunct Professor, Department of Obstetrics and Gynaecology, National University of Singapore and Director, Embryonics International, Gleneagles Hospital, Singapore for his time to time advice and encouragements. Dr. S. L. Liow, Chief embryologist and lab manager, Embryonics International, Singapore for his advice and help in ICSI and related techniques. Mrs. Ruhaima Ramli, Embryologist, IVF Unit, Pantai Medical Centre, Kuala Lumpur, Malaysia for initially helping me with ICSI technique. Mrs. Nooraain Hashim, Lecturer, Faculty of Science, Mara University of Technology, Shah Alam, Selangor, Malaysia for helping in embryo staining and fluorescence microscopy.

I am grateful to Department of Veterinary Services, Malaysia for providing me with the access to their abattoirs and Mr. Gandhi of Abattoir Complex, Shah Alam, Selangor for his constant help in getting the goat ovaries.

The 'OR Team' members are highly acknowledged for their constant help during LOPU.

Special thanks should go to Ms. Nor Fadillah for being helpful all the time, Ms. Raja Ili for helping in pregnancy diagnosis and Ms. Phek Jin for her special help.

I give thanks with all my heart the present and ex-members of ABEL, NaTuRe and ISB Mini Farm. I specially would like to recognise the following people: Parani Baya, Sow Chan, Mohd. Noor Hisham, Alice Phua, Shariffah Nazari, Izuan Bahtiar, Anna Amir, Ainul Bahiyah, Wai Kit, Razali Jonit, Asmad Kari, Nor Azman, Li Ling, Suzuna Baharuldin, Jamaludin Alias, Raman Kutty, Hazlan Che Haroon, Erma Farina,

Ravindran, Nor Fazliana and Farid. I consider myself a very lucky person for having met and shared some unforgettable time with all of you.

This work was supported by grants from MOSTI Special Project (grant no. 01-02-03-0696) and IPPP (grant nos. Vote F-0179/2004D, 0145/2005D and P0170/2006C). I acknowledge these financial help without which this study would not be possible.

I am grateful and would like to thank the Institute of Biological Sciences, Faculty of Science as well as the University of Malaya for accepting me as a PhD candidate.

I am very thankful to Islamic Development Bank (IDB) for awarding me 'IDB Merit Scholarship for Higher Technology' for pursuing my PhD study.

I am also thankful to Bangladesh Open University for granting me the study leave required to peruse the PhD study in the University of Malaya.

I would like to thank my mother, elder brother, sisters, other family members and friends for their encouragement during the course of my study.

I would like to recognize my wife and two kids for all the sacrifices that they have made to enable me to pursue an advance degree.

Finally, I would like to remember my late father who always encouraged me for higher study.